


ORI-02

**DE BEERS**  
A DIAMOND IS FOREVER

STRICTLY CONFIDENTIAL

**Closing Report on Reconnaissance Permit  
7476/IV (GD) SM 3/2000 Orissa**

Report for the period  
16/02/05 to 15/08/07



*In terms of the relevant legislation, the information reported in this document is to be kept strictly confidential by the orissa State Government for the period of two years from the date of expiry of the license.*



**DE BEERS INDIA PRIVATE LIMITED**

Advance Business Centre, No 83, Maker Chambers IV, Nariman Point, Mumbai 400021  
Tel: 022-22832927 Fax: 022-22832823

**Closing Report on Reconnaissance Permit  
7476/IV (GD) SM 3/2000 Orissa**

Report for the period  
16/02/05 to 15/08/07

**1. Reconnaissance Permit (RP) Status**

The RP is 1880 Sq. Km in extent and was executed at Bhubaneswar on 15<sup>th</sup> February 2005. As per rule 7(i) (a) of MCR 1960, it is scheduled to be reduced by 50% on or before 14<sup>th</sup> February 2007. A total of 1330 Sq. Km was relinquished on 20<sup>th</sup> July 2006 and subsequent to this the remaining 550 Sq. Km was also relinquished on 15<sup>th</sup> August 2007. This report summarizes the exploration work carried out in the total area of 1880 sq km till 15<sup>th</sup> August 2007 (Map 1).

**2. Geology and Geomorphology of the RP area**

The Bhandara Craton is rectangular in shape and bounded by the Mahanadi Rift in NE, the Godavari Rift in SW, Narmada Rift to the NW, and the NE-SW trending Eastern Ghat Mobile Belt on the SE.

The Eastern Ghat Mobile Belt is juxtaposed with the Archean craton along a ductile thrust with a northwesterly thrust movement. This shear zone is characterized by south-easterly dipping mylonitic foliation and a down-dip stretching lineation (the L-S fabric of mylonites).

Geologically the RP area consists of the eastern margin of the Bhandara Craton with a complex lithological association of basic intrusive rocks, granitoids, supracrustal rocks and unmetamorphosed Late Proterozoic sediments in the west and the Eastern Ghat Mobile Belt in the eastern side with high grade metamorphic rocks (Map 2).

Uncertainty and unavailability of precise age data of the rocks have resulted in varied interpretations and correlations attempted by different workers. A simplified stratigraphic succession of the Bhandara Craton is presented below. The Sukma and Bengpal metamorphic rocks are possibly the oldest Mesoarchaeon sequences of this region. The relationship of the precursors of the trondhjemitic gneisses (3.56 Ga) with the Sukma and Bengpal sequence is debatable from being a probable intrusive to being basement.

The Sukma supracrustal rocks consists of high grade metamorphosed orthoquartzites, carbonates and pelites in association with prominent iron formations and subordinate mafic ultramafic rocks. The Bengpal Group overlies the Sukma metamorphics with an unconformity and consists of amygdular metabasalts and andalusite-chlorite schist which are overlain by conglomerate and quartzites.

However, Ramakrishnan (1990) assigns Palaeoproterozoic ages (2.5Ga and 2.3Ga) for the Sukma metamorphics and Bengpal Group.



DE BEERS INDIA PRIVATE LIMITED

Advance Business Centre, No 83, Maker Chambers IV, Nariman Point, Mumbai 400021  
Tel: 022-22832927 Fax: 022-22832823

The Bailadila Group overlies the Bengpal Group with an unconformity and is composed of feldspathic quartzites, chloritic phyllite and banded iron formations.

The Eastern Ghat Mobile Belt is recognised as an accreted terrain with high-grade metamorphic rocks. The major mappable units are mafic schists, quartz-feldspar-hypersthene granulites (charnockites), sillimanite-garnet-cordierite-sapphirine-schist and gneisses (metasediments locally known as khondalites) also the Calc-silicates, mafic granulites, leptynites (leucocratic quartz-feldspar-garnet gneiss), intrusions of anorthosite, gabbro, norite, dunite and serpentinite with chromite, and alkaline rocks. Most of these rock suites have been subjected to three or four deformational events and with a large number of shear zones cutting across them, and together comprise the Khariar-Angul Metapelite Complex of the Sukma Group.

Late Proterozoic detached basins containing unmetamorphosed sediments overlie the Archaean rocks of the Bhandara Craton. There are two major and six minor basins in the Craton. Two smaller basins, the Neogarh-Khariar (Nawapara-Kharihar) and the Ampani Basins are located within the RP area.

The Khariar Basin is composed of an alternating sequence of sandstones and shale and is equivalent to the Chandrapur Group of the Chhattisgarh Basin.

The Ampani Basin forms an outlier south of the Khariar Basin and is composed principally of conglomerates, sandstones, siltstones, purple shales and calcareous intercalations. The rock association resembles the Chandrapur Group of the Chhattisgarh Basin and the Tirahgarh Formation of the Indravati Basin.

Simplified stratigraphic succession of the Bhandara Craton

Chhattisgarh Supergroup	Un-metamorphosed Sediments	Upper Proterozoic
~~~~~	Unconformity	~~~~~
Basic Dykes	Dolerites, Amphibolites	
~~~~~	Unconformity	~~~~~
Bailadila Group	Banded Iron Formation, Phyllites with chert and local conglomerates	ca. 2.9-2.5 (Bandhopadhyay et al 1995). ca. 2.1 Ga (Ramkrishnan 1990)
~~~~~	Unconformity	~~~~~
Bengpal Group	Amygdular metabasalt, andalusite schist and quartzite	ca. 2.9-3.5 (Bandhopadhyay et al 1995). 2.3 Ga (Ramkrishnan 1990)
~~~~~	Unconformity	~~~~~
Sukma (Group ?) metamorphic suite	Sillimanite quartzite, cordierite anthophyllite rocks, Banded Iron Formation	ca. 2.9-3.5 (Bandhopadhyay et al 1995). ca. 2.5 Ga (Ramkrishnan 1990)
~~~~~	Unconformity	~~~~~
	Basement Gneiss (TTG)?	3.56 (?): Zircons of tonalites by U-Pb method.



**DE BEERS INDIA PRIVATE LIMITED**

Advance Business Centre, No 83, Maker Chambers IV, Nariman Point, Mumbai 400021  
Tel: 022-22832927 Fax: 022-22832823

The RP area is within the drainage basin of the Sundar Nadi which joins with Tel River (a tributary of the River Mahanadi) at the SE corner of the RP (Map 3). The western part of the RP comprise elevated plateaus rising to over 700m. Below is a gently sloping plain falling from approximately 250m in the northwest, to 180m in the southeast. Isolated hills occur in the north and NE rise to over 700m above the plain.

The western plateau comprises sandstones of the Khariar Basin, whilst the plains are underlain by metapelites of the Sukama Group.

### **3. Activity during the period (16/02/05 to 15/08/07)**

#### **3.1 Reconnaissance Sampling**

A total of 169 reconnaissance stream/loam samples were collected from the RP.

Stream samples comprised 75 litres of unscreened material, collected from natural heavy mineral trapsites and field screened to -2.0mm.

Loam samples comprise 30 litres of material collected from surface in interfluves.

Reconnaissance sample localities and information are shown in Map 4 and Table 1.

The samples were processed at De Beers heavy mineral treatment plant in Bangalore, and the concentrates were consigned to the Bangalore Mineralogical Lab, India for further processing and sorting.

#### **3.2 Reconnaissance Sampling Results**

Received results for most of the samples and 45 samples reported positive with respect to kimberlitic indicator minerals (Map 5 and Table 2). A Total of 192 spinels and 2 ilmenites were reported.

#### **3.3 Airborne Hyperspectral Scanner Survey**

An airborne survey utilizing De Beers's proprietary Hyper Spectral Scanner was completed during March 2005 (Map 6). The system works by measuring reflectance of narrow wavelength bands of sunlight reflected from the Earth's surface. Different minerals (as well as other materials) absorb different wavelengths of light to varying degrees. The Airborne Hyperspectral scanner Survey system is sensitive enough to actually distinguish some specific types of minerals by the absorption of certain wavelengths of light detected. In the search for kimberlites, the system is configured to look for the presence of magnesium-rich clay minerals, derived from the weathering of ultramafic rocks.

There were 32 anomalies selected for field follow-up (Map 7).



DE BEERS INDIA PRIVATE LIMITED

Advance Business Centre, No 83, Maker Chambers IV, Nariman Point, Mumbai 400021  
Tel: 022-22832927 Fax: 022-22832823

Follow up of the survey involved field visits of anomalies and identification of the causative lithological units to conclude whether the source of the anomaly is kimberlitic or not. All the anomalies were visited (Table 3 and Map 8). Samples were collected from surface material for PIMA (Portable Infra-Red Mineral Analyses) to confirm the causative material.

Detailed sheets of Airborne Hyper spectral scanner survey anomalies followed up are attached as Appendix 1

### **3.4 Airborne Geophysical Survey**

The Airborne Geophysical survey data were acquired from DMG Orissa for the RP area. A Fixed wing Airborne Magnetic and Time Domain EM survey was also carried out by Fugro over part of the RP area as Sinapali Block. The results were processed and interpreted.

Based on the response, a total of 34 Aeromagnetic anomalies were identified (Map 9) and followed up with ground geophysical surveys, wherever necessary.

### **3.5 Ground Geophysical Survey**

#### **3.5.1 Ground Magnetic Survey**

Sixteen magnetic anomalies were followed up with ground magnetic survey using a Proton precision magnetometer (Table 4), covering a total of 188 line kilometres (Map 10).

Detailed sheets of Magnetic anomalies followed up with ground survey are attached as Appendix 2

#### **3.6 Drilling**

A total of 201 meters in 16 holes were drilled to test the geophysical anomalies and Airborne Hyper Spectral anomalies (Table 5 and Map 11).

Detailed sheets of borehole logs are attached as Appendix 3

## **4. Interpretation**

Based on the results received for various techniques used in exploration of this RP, the western portion has been identified as having the most potential for the discovery of a diamond bearing body (Fig 1 – 10). This area was retained in order to carry out further exploration. However, further exploration work in this area failed to establish the presence of a significantly diamond rich kimberlite/ related rock and hence this RP is completely relinquished (Map 1).



DE BEERS INDIA PRIVATE LIMITED

Advance Business Centre, No 83, Maker Chambers IV, Nariman Point, Mumbai 400021

Tel: 022-22832927 Fax: 022-22832823

**5. Personnel**

<b>Name</b>	<b>Designation</b>	<b>Education</b>
K.V.Suryanarayana Rao	Project Manager	M.Sc. Tech-Applied Geology
Krishna Pande	Staff Geologist	M.Sc. – Applied Geology
Chandan kumar	Staff Geologist	M.Sc. – Applied Geology
K.V.Praveen Kumar	Staff Geologist	M.Tech-Remote Sensing
Prashant Laharia	Geologist (on contract)	M.Sc Tech Geology
Sukhbinder Sharma	Geologist (on contract)	M.Sc Geology
Sudipta Sarkar	Geologist (on contract)	M.Sc. Tech-Geology
Rekha K.R.	Kimberlitic Mineral Analyst	M.Sc Geology
Shobha N.	Kimberlitic Mineral Analyst	M.Sc. Geology
Sanjay Deogiri	IT Manager	B.Sc. Electronics, MCSE
Rina David	SHE Officer	Graduate
Girish Menon	Security Adviser	Grade 12
Gajanana Naik	Treatment Plant Supervisor	Graduate
R.Shrinivasalu	Field Assistant	Grade 10
Venu Kumar	Field Driver	Grade 12
A.Channaiah	Field Driver	Grade 10
Rajanna	Field Driver(on contract)	Grade 10

**Labour**

Labourers were employed on a daily basis from local towns and villages to help with the field work.

**6. Training**

De Beers maintains high operating standards including safety and environmental awareness. To this end, training is an integral part of career development with the organization. The following is a short summary of training completed during reporting period.

All Earth scientists, field assistants and field drivers attended a training program on Crisis Management to deal emergency situation related to safety and security in the field.

An HIV/AIDS awareness programme was conducted for the entire field based staff.

The drivers attended a training course on Defensive Driving and Road Safety conducted by the Automobile Association of Southern India.

All field drivers attended training on vehicle maintenance conducted by Mahindra & Mahindra.

Earth Scientists attended training on new mineral classification scheme.



**DE BEERS INDIA PRIVATE LIMITED**

Advance Business Centre, No 83, Maker Chambers IV, Nariman Point, Mumbai 400021  
Tel: 022-22832927 Fax: 022-22832823

All field geologists attended a training program on First Aid conducted by International SOS.

Earth scientists attended a programme on Arc SDE and Arc GIS to hone their skills in Database management.

Earth Scientists also attended a programme on Datamine solutions which will enable them to effectively organize data from drilling programmes.

A brief introductory lecture on basic geology was given to field drivers and field assistants.

Earth Scientists attended a programme on XSIMS database management.

Earth Scientist completed a sort time assignment for drilling and petrology training in Canada and South Africa.

### **7. Expenditure**

Total expenditure of Rs. 36,297,818 has been incurred for the Reconnaissance Permit to date. The expenditure was incurred as follows:


Capital expenditure: Rs. 3,598,071

Revenue expenditure: Rs. 32,699,747

### **8. References**

Bandyopadhyay, B. K. Roy A. Jain A. K. 1995: Structure and Tectonics of a part of the Central Indian Shield: In Continental crust of north-western and central India. Sinha Roy S, Gupta K. R. (Eds) Geol Soc India, Mem 31, 433-468.

Ramakrishnan M 1990, Crustal development in southern Bastar, Central Indian Craton. In Precambrian of central India Geol, Surv, India Spl Publ. No 28 44-66

  
KVSuryanarayana Rao  
Technical Specialist - Geology  
De Beers India Private Ltd



**DE BEERS INDIA PRIVATE LIMITED**

Advance Business Centre, No 83, Maker Chambers IV, Nariman Point, Mumbai 400021  
Tel: 022-22832927 Fax: 022-22832823